Markovian N-Server Queues

(Birth & Death Models)



Arrivals Poisson (λ) ;

Services exp(μ) $(E(S) = 1/\mu)$

Servers N **statistically identical**, serving FCFS.

Offered load $R = \lambda \times E(S) = \lambda/\mu$ Erlangs

$Q(t)$ = number in system (served + queued) at time t.

$W(k)$ = queueing time of k-th arrival.

Steady State: $Q(\infty)$, $W(\infty)$, when exists.

non-idling $\Rightarrow$ $[Q(t) - N]^+$ = queue-length, and

$[Q(t) - N]^-$ = number of idle servers

Examples: M/M/N/K, Abandonment, Balking

M/M/∞: $\lambda_k \equiv \lambda, \quad \mu_k = k \cdot \mu, \qquad k \geq 1.$

Steady state $\pi_k = e^{-R} R^k / k!$ Poisson (R)

M/M/N/N: $\lambda_k \equiv \lambda, \quad \mu_k = k \cdot \mu, \quad 0 \leq k \leq N.$

Steady state $$\pi_k = \frac{R^k}{k!} \Big/ \sum_{n=0}^{N} \frac{R^n}{n!}$$

Erlang-B $P\{\text{Blocked}\} \doteq \boldsymbol{E_{1,N}} = \pi_N$, by PASTA

M/M/N: $\lambda_k \equiv \lambda, \qquad \mu_k = (k \wedge N) \cdot \mu \quad , \quad k \geq 1.$

Steady state $\Leftrightarrow \ \rho = \dfrac{\lambda}{N\mu} < 1$, servers' utilization.

Erlang-C $P\{\text{Wait} > 0\} \doteq \boldsymbol{E_{2,N}} = \displaystyle\sum_{k \geq N} \pi_k$

$$W(\infty) \mid W(\infty) > 0 \ \stackrel{d}{=} \ \exp\left(\text{mean} = \frac{1}{N\mu(1-\rho)}\right)$$

Restriction to a Set via Time-Change

X Markov , $X(\infty) \sim \pi$

$\tau_A(t) = \int_0^t 1_{\{X(u) \in A\}} du$ time in A



$X_A(t) = X\left(\tau_A^{-1}(t)\right)$ X restricted to A: Markov

$X_A(\infty) \sim X(\infty) | X(\infty) \in A$

Example: M/M/N/N $\stackrel{d}{=}$ M/M/∞ restricted to $\{0, 1, \ldots, N\}$.

$$E_{1,N} = Pr\{X_R = N\}/Pr\{X_R \leq N\},$$

$$X_R \sim \text{Poisson}\,(R)$$

Example: M/M/N (Erlang-C)



$Q(t) =$ number in system at time $t \geq 0$

$Q_- = Q$ restricted to $\{0, 1, \ldots, N-1\}$: M/M/N-1/N-1 ; λ, μ.

$Q_+ = Q$ restricted to $\{N, N+1, \ldots\}$: M/M/1 ; $\lambda, N\mu$.

Evolution of Q: Alternates between M/M/1 (Q_+) and M/M/N-1/N-1 (Q_-).

$$P(\text{Wait} > 0) = \boldsymbol{E_{2,N}} = \frac{T_{N,N-1}}{T_{N,N-1} + T_{N-1,N}} \quad , \quad \text{by PASTA,}$$

where (after conditioning on the first step):

$$T_{N,N-1} = \frac{1}{\mu_N \pi_+(0)} = \frac{1}{N\mu(1-\rho)} \quad ; \quad \rho = \frac{\lambda}{N\mu}$$

$$T_{N-1,N} = \frac{1}{\lambda_{N-1} \pi_-(N-1)} = \frac{1}{\lambda\, E_{1,N-1}} \quad .$$

Hence,

$$E_{2,N} = \left[1 + \frac{T_{N-1,N}}{T_{N,N-1}}\right]^{-1} = \left[1 + \frac{1-\rho}{\rho\, E_{1,N-1}}\right]^{-1},$$

in which

$$\boldsymbol{E_{1,N}} = \frac{R^N}{N!} \Big/ \sum_{k=0}^{N} \frac{R^k}{k!} \quad ; \quad R = N\rho = \tfrac{\lambda}{\mu}$$

M/M/N/N (Erlang-B) with Many Servers: $\mathrm{N} \uparrow \infty$

Assume: μ fixed, while $\lambda_N \uparrow \infty$ as $N \uparrow \infty$.

Recall: $R = R_N = \lambda_N/\mu, \quad \rho = \rho_N = R_N/N.$

Erlang-B: $E_{1,N} = Pr\{X_R = N\}/Pr\{X_R \le N\},$

where $X_R \stackrel{d}{=}$ Poisson (R).

R large $X_R \stackrel{d}{\approx} R + Z\sqrt{R}$, where $Z \stackrel{d}{=} N(0,1)$,

suggesting $N \sim R + \beta\sqrt{R}, \ -\infty < \beta < \infty.$

Then $E_{1,N} = Pr\{N-1 < X_R \le N\}/Pr\{X_R \le N\}$

$$\sim Pr\left\{\beta - \tfrac{1}{\sqrt{R}} < Z \le \beta\right\}/Pr\{Z \le \beta\}$$

$$\sim \tfrac{1}{\sqrt{R}}\,\varphi(\beta)/\phi(\beta) \approx \tfrac{1}{\sqrt{N}}\,\tfrac{\varphi(\beta)}{\phi(\beta)} =$$

$$= \tfrac{1}{\sqrt{N}}\,\tfrac{\varphi(-\beta)}{1-\phi(-\beta)} = \tfrac{1}{\sqrt{N}}\,h(-\beta), \quad h = \text{hazard rate.}$$

Algebra: μ fixed, $N \uparrow \infty$.

QED: $N \sim R + \beta\sqrt{R}$, for some $\beta, -\infty < \beta < \infty$

$$\Leftrightarrow \lambda_N \sim \mu N - \beta\mu\sqrt{N}$$

$$\Leftrightarrow \rho_N \sim 1 - \frac{\beta}{\sqrt{N}}, \quad \text{namely} \quad \lim_{N\to\infty} \sqrt{N}\,(1-\rho_N) = \beta.$$

Theorem: (Jagerman, 1974) QED $\Leftrightarrow \lim_{N\to\infty} \sqrt{N}\,E_{1,N} = h(-\beta)$.

M/M/N (Erlang-C) with Many Servers: $\mathrm{N} \uparrow \infty$



$Q(0) = N$: all servers busy, no queue.

Recall $$E_{2,N} = \left[1 + \frac{T_{N-1,N}}{T_{N,N-1}}\right]^{-1} = \left[1 + \frac{1-\rho_N}{\rho_N E_{1,N-1}}\right]^{-1}.$$

Here $$T_{N-1,N} = \frac{1}{\lambda_N E_{1,N-1}} \sim \frac{1}{N\mu \times h(-\beta)/\sqrt{N}} \sim \frac{1/\mu}{h(-\beta)\sqrt{N}}$$

which applies as $\sqrt{N}\,(1-\rho_N) \to \beta,\ -\infty < \beta < \infty.$

Also $$T_{N,N-1} = \frac{1}{N\mu(1-\rho_N)} \sim \frac{1/\mu}{\beta\sqrt{N}}$$

which applies as above, but for $0 < \beta < \infty$.

Hence, $$E_{2,N} \sim \left[1 + \frac{\beta}{h(-\beta)}\right]^{-1},$$ assuming $\beta > 0$.

QED:
$$N \sim R + \beta\sqrt{R} \quad \text{for some} \quad \beta, \quad 0 < \beta < \infty$$
$$\Leftrightarrow \quad \lambda_N \sim \mu N - \beta\mu\sqrt{N}$$
$$\Leftrightarrow \quad \rho_N \sim 1 - \frac{\beta}{\sqrt{N}}, \quad \text{namely} \quad \lim_{N\to\infty} \sqrt{N}\,(1-\rho_N) = \beta.$$

Theorem (Halfin-Whitt, 1981) QED $\Leftrightarrow \lim_{N\to\infty} E_{2,N} = \left[1 + \frac{\beta}{h(-\beta)}\right]^{-1}$.

QED M/M/N: Steady-State

Theorem (Halfin-Whitt, 1981)

Consider a sequence M/M/N, N = 1,2,.... Then the following are equivalent

- $\lim_{N\to\infty} P\{W_N(\infty) > 0\} = \alpha$, for some $0 < \alpha < 1$;
- $\lim_{N\to\infty} \sqrt{N}\,(1-\rho_N) = \beta$, for some $0 < \beta < \infty$;
- $\lambda_N = \mu N - \mu\beta\sqrt{N} + o\left(\sqrt{N}\right)$, i.e., $N \sim R_N + \beta\sqrt{R_N}$,

in which case

- $\alpha = \alpha(\beta) = \left[1 + \dfrac{\beta}{h(-\beta)}\right]^{-1}$ Halfin-Whitt function
- $\sqrt{N}\,W_N(\infty) \stackrel{d}{\to} \hat{W}(\infty)$, $\hat{W}(\infty) \mid \hat{W}(\infty) > 0 \stackrel{d}{=} \exp(\mu\beta)$.

Moreover (Queue must be order $\sqrt{N}$)

$$\left(\frac{1}{\sqrt{N}}\,[Q_N(\infty) - N]^+, \sqrt{N}\,W_N(\infty)\right) \stackrel{d}{\to} \left(\hat{Q}^+(\infty),\, \hat{W}(\infty)\right)$$

where $\hat{Q}^+(\infty) = \mu\hat{W}(\infty) \stackrel{d}{=} \exp(\beta)$.

Proof: Let $\wedge_N(\cdot)$ be Poisson(λ_N). Then divide by $\sqrt{N}$ and take $N \uparrow \infty$ in

$$\wedge_N[W_N(\infty)] \stackrel{d}{=} [Q_N(\infty) - N]^+ \text{ (Haji+Newell, 1971).}$$

QED M/M/N : Process View

Framework : Sequence of M/M/N systems,

indexed by $N = 1, 2, \ldots$

- $Q_N = \{Q_N(t),\ t \geq 0\}$ number in system
- $V_N = \{V_N(t),\ t \geq 0\}$ virtual waiting time, under FCFS

Parameters	$\lambda_N\ ,\ \mu_N \equiv \mu$
Offered load	$R_N = \lambda_N \times \frac{1}{\mu} = \lambda_N/\mu$
Traffic intensity	$\rho_N = R_N/N$

Each M/M/N is a Birth & Death process, which is ergodic iff $\rho_N < 1$, in which case $\rho_N =$ servers' utilization.

QED Scaling : $\lambda_N = N\mu - \beta\mu\sqrt{N}$, namely

$$\rho_N = 1 - \frac{\beta}{\sqrt{N}}$$

Approximations of Process and Stationary Distribution, as $N \uparrow \infty$:

- $\widehat{Q}_N(t) = \frac{1}{\sqrt{N}}[Q_N(t) - N]\ , \quad 0 \leq t \leq \infty$
- $\widehat{V}_N(t) = \sqrt{N} V_N(t).$

Number in System, Centered and Rescaled: $\hat{Q}_N(t)$

$$(\mu = 1, \beta = 0.5)$$









Thanks to G. Shaikhet for the simulations and insight.

Approximating Queueing and Waiting

- $Q_N = \{Q_N(t),\ t \geq 0\}$: $Q_N(t) =$ number in system at $t \geq 0$.
- $\hat{Q}_N = \{\hat{Q}_N(t), t \geq 0\}$: stochastic process obtained by centering and rescaling:

$$\hat{Q}_N = \frac{Q_N - N}{\sqrt{N}}$$

- $\hat{Q}_N(\infty)$: stationary distribution of $\hat{Q}_N$
- $\hat{Q} = \{\hat{Q}(t), t \geq 0\}$: process defined by: $\hat{Q}_N(t) \xrightarrow{d} \hat{Q}(t)$.



Approximating (Virtual) Waiting Time

$$\hat{V}_N = \sqrt{N}\, V_N \Rightarrow \hat{V} = \left[\frac{1}{\mu}\,\hat{Q}\right]^+ \qquad \text{(Puhalskii, 1994)}$$

Diffusion Processes in $\mathbb{R}^1$

$X = \{X_t,\ t \geq 0\}$ Markov process with continuous sample paths

Kolmogorov/Feller/Dynkin: characterized (Strong Markov, Continuous) by

- Drift function $\mu_t(x)$ (infinitesimal mean)

$$E\left[X_{t+\epsilon} - X_t \mid X_t = x\right] = \mu_t(x)\epsilon + o(\epsilon), \quad \epsilon \downarrow 0;$$

- Diffusion function $\sigma_t(x)$ (infinitesimal variance)

$$\text{Var}\left[X_{t+\epsilon} - X_t \mid X_t = x\right] = \sigma_t(x)\epsilon + o(\epsilon), \quad \epsilon \downarrow 0;$$

- Boundary behavior: **inaccessible**; absorbing, sticky, **reflecting**

Time-homogeneous: $\mu_t(x) \equiv \mu(x), \qquad \sigma_t(x) \equiv \sigma(x).$

Examples:

1. $\mu(x) \equiv \mu\ ;\ \sigma(x) \equiv \sigma > 0$ Brownian Motion: BM (μ, σ^2)

   $\mu = 0 \qquad \sigma = 1$ Standard BM (SBM)



Diffusions: Examples (Continued)

2. $\mu(x) = a - bx,\ \sigma(x) \equiv \sigma$ Orenstein-Uhlenbeck (OU)

   Stationary distribution $\Leftrightarrow\ b > 0 : N(\frac{a}{b}, \frac{\sigma^2}{2b})$

3. Reflected BM (μ, σ^2), reflected at 0. (Harrison's book)

$$\begin{cases} dX_t = \mu dt + \sigma dB_t + dY_t, \quad X_0 \geq 0 \\ X \geq 0,\ dY \geq 0,\ XdY = 0 \end{cases}$$

   Stationary distribution $\Leftrightarrow \mu < 0 :\ \exp(2|\mu|/\sigma^2)$

4. Reflected OU on $[0, \infty)$, (Glynn and Ward), or generally (Dupuis)

$$\begin{cases} dX_t = \mu_t(X_t)dt + \sigma_t(X_t)dB_t + dY_t; \quad X_0 \geq 0 \text{ given} \\ X \geq 0,\ dY \geq 0,\ XdY = 0 \end{cases}$$

Ito: Characterized by solutions to stochastic differential equations

$$dX_t = \mu_t(X_t)dt + \sigma_t(X_t)dB_t, \quad t \geq 0; \quad X_0 \text{ given.}$$

(Reference: Karlin and Taylor, "2nd Course"; Karatzas and Shreve.)

Formalizes the "infinitesimal description":

$$X_{t+\epsilon} - X_t \mid X_t = x \overset{d}{\approx} \mu_t(x) \cdot \epsilon + \sigma_t(x) \cdot (B_{t+\epsilon} - B_t)$$

where $B_{t+\epsilon} - B_t \sim N(0, \epsilon)$, independent of $\{B_u,\ u \leq t\}$.

(Equivalently, $B = \{B_t,\ t \geq 0\}$ is SBM.)

QED M/M/N : Diffusion Approximation

$$
\begin{aligned}
Q_N &= \{Q_N(t),\ t \geq 0\} && \text{Birth and Death with} \\
\lambda_N &= \mu N - \mu\beta\sqrt{N} && \text{birth rate, constant} \\
\mu_N(x) &= \mu \cdot (x \wedge N) && \text{death rate at state } x \\
\hat{Q}_N(t) &= \frac{1}{\sqrt{N}}\left[Q_N(t) - N\right] && \text{Scaled queue length of} \\
& && \text{customers } (+) \text{ or servers } (-)
\end{aligned}
$$

Diffusion: $\text{Var}\left[\hat{Q}_N(t+\epsilon) - \hat{Q}_N(t) \mid \hat{Q}_N(t) = x\right] = 2\mu \cdot \epsilon + o(\epsilon)$

Drift: $E\left[\hat{Q}_N(t+\epsilon) + \hat{Q}_N(t) \mid \hat{Q}_N(t) = x\right] =$

$$
\frac{1}{\sqrt{N}}\ E\left[Q_N(t+\epsilon) - Q_N(t) \mid Q_N(t) = N + x\sqrt{N}\ \right] =
$$

$$
\frac{1}{\sqrt{N}}\ \left[\lambda_N - \mu_N\left(N + x\sqrt{N}\ \right)\right] \cdot \epsilon + o(\epsilon) =
$$

$$
\frac{1}{\sqrt{N}}\ \left\{\mu N - \mu\beta\sqrt{N}\ - \mu\left[\left(N + x\sqrt{N}\ \right) \wedge N\right]\right\} \cdot \epsilon + o(\epsilon) =
$$

$$
\left.\begin{array}{lll}
= & -\mu\beta \cdot \epsilon + o(\epsilon) & x \geq 0 \quad \text{(BM)} \\
= & -\mu(\beta + x) \cdot \epsilon + o(\epsilon) & x \leq 0 \quad \text{(OU)}
\end{array}\right\} = -\mu(\beta - x^-) \cdot \epsilon + o(\epsilon)
$$

Expect $\hat{Q}_N \xrightarrow{d} \hat{Q}$ diffusion: $\mu(x) = -\mu(\beta - x^-),\ \sigma(x) = 2\mu$

Proof: Apply Stone (1963), as in Halfin-Whitt (1981).

QED M/M/N : Diffusion Approximation

Theorem (Halfin-Whitt).

Consider a sequence M/M/N, N = 1,2,. . .. Assume QED(β).

Define $\hat{Q}_N(t) = \frac{1}{\sqrt{N}}\,[Q_N(t) - N]$, $0 \le t < \infty$,

$\hat{V}_N(t) = \sqrt{N}\, V_N(t)$, $0 \le t < \infty$.

If $\left(\hat{Q}_N(0),\ \hat{V}_N(0)\right) \xrightarrow{d} \left(\hat{Q}(0),\ \hat{V}(0)\right)$, then

$$\left(\hat{Q}_N,\ \hat{V}_N\right) \xrightarrow{d} \left(\hat{Q},\ \frac{1}{\mu}\,\hat{Q}^+\right)$$ (Functional CLT),

where $\hat{Q}$ is a diffusion process starting at $\hat{Q}(0)$, with infinitesimal parameters

$$\mu(x) = \begin{cases} -\mu\beta & x \ge 0 \\ -\mu(x+\beta) & x < 0 \end{cases} \quad , \quad \sigma^2(x) = 2\mu \ ;$$

and steady-state distribution $\hat{Q}(\infty)$ given by

$P\{\hat{Q}(\infty) \ge 0\} = \alpha(\beta)$,

$P\{\hat{Q}(\infty) > x \mid \hat{Q}(\infty) \ge 0\} = e^{-x\beta}$ (exp $\leftrightarrow$ RBM)

$P\{\hat{Q}(\infty) \le x \mid \hat{Q}(\infty) \le 0\} = \phi(x+\beta)/\phi(\beta)$ (normal $\leftrightarrow$ OU)

E-Driven M/M/N : Approximations

Recall: 1:1 staffing gave rise to $\lambda_N \sim \mu N - \mu\gamma$

$$\lim_{N\to\infty} N(1-\rho_N) = \gamma, \quad 0 < \gamma < \infty.$$

Applying the usual QED scaling: $\hat{Q}_N(t) = \frac{1}{\sqrt{N}}(Q_N(t) - N)$,

results in a Halfin-Whitt diffusion limit with $\beta = 0$: No stationary distribution for

$$\hat{Q}^+(t) = \lim_{N\to\infty} \frac{1}{\sqrt{N}}(Q_N(t) - N)^+.$$

To identify a more informative scaling, recall:

$$W_N(\infty) \mid W_N(\infty) > 0 \stackrel{d}{=} \exp\left(\text{mean} = \frac{1}{N\mu(1-\rho_N)}\right).$$

Also,

$$\lim_{N\to\infty} P\{W_N(\infty) > 0\} = 1.$$

Hence,

$$W_N(\infty) \stackrel{d}{\to} \hat{W}(\infty) \stackrel{d}{=} \exp(\mu\gamma).$$

And (by Haji & Newell)

$$\tfrac{1}{N}\,[Q_N(\infty) - N]^+ \stackrel{d}{\to} \hat{Q}(\infty)^+ \stackrel{d}{=} \mu\hat{W}(\infty) \stackrel{d}{=} \exp(\gamma)\,.$$

Expect a non-degenerate behavior of

$$\bar{Q}_N(t) = \frac{1}{N}[Q_N(t) - N] \stackrel{d}{\to} \bar{Q}(t) \;\;,\; 0 \le t \le \infty$$

Process View: Simulations of $\bar{Q}_N(t)$



Proposition: Let $\lim_{N\to\infty} N(1-\rho_N) = \gamma, \quad 0 < \gamma < \infty.$

If $\frac{1}{N}[Q_N(0) - N] \xrightarrow{d} \bar{Q}(0)$, then

$$\bar{Q}_N(t) \xrightarrow{d} \bar{Q}(t) = \begin{cases} \bar{Q}(0) & , \quad \text{if } \bar{Q}(0) \geq 0, \\ \bar{Q}(0)e^{-\mu t} & , \quad \text{if } \bar{Q}(0) \leq 0. \end{cases}$$

Proof:

Diffusion: $\text{Var}\left[\bar{Q}_N(t+\epsilon) - \bar{Q}_N(t) \mid \bar{Q}_N(t) = x\right] = 0 + o(\epsilon)$

Drift: $E\left[\bar{Q}_N(t+\epsilon) + \bar{Q}_N(t) \mid \bar{Q}_N(t) = x\right] =$

$$\left.\begin{array}{rl} = & 0 \cdot \epsilon + o(\epsilon) \quad x \geq 0 \\ = & -\mu x \cdot \epsilon + o(\epsilon) \quad x \leq 0 \end{array}\right\} = \mu x^- \cdot \epsilon + o(\epsilon).$$

Conclude $\bar{Q}_N \xrightarrow{d} \bar{Q}$: $\mu(x) = \mu x^-, \ \sigma(x) \equiv 0$.

Discovered that $\bar{Q}_N(t), 0 \leq t < \infty$ degenerates, as $N \uparrow \infty$. But

$$\bar{Q}_N(\infty) = \frac{1}{N}[Q_N(\infty) - N]^+ \xrightarrow{d} \exp(\gamma). \qquad \textbf{What's going on?}$$

Two explanations (assuming $\bar{Q}(0) \geq 0$, non-random, for simplicity):

1. For $k \geq N$ (with $\lambda = N\mu - \gamma\mu$),

$$T_{k+cN,k} = \theta(N) * T_{k+1,k} = \theta(N) * \theta(1) \to \infty, \quad \text{as } N \uparrow \infty.$$

Hence, $\bar{Q}_N(t)$ takes close to infinite time to move a one-unit distance, suggesting that, in the limit, $\bar{Q}(t)$ freezes.

2. Consider Q_N^+: the restriction of the Q_N to $\{N, N+1, ..\}$.

$Q_N^+ \stackrel{d}{=} M/M/1$ with $\lambda^+ = N\mu - \gamma\mu$, $\mu^+ = N\mu$.

$\stackrel{d}{=} M/M/1$ with $\bar{\lambda} = \frac{\lambda^+}{N} \to \mu$, $\bar{\mu} = \frac{\mu^+}{N} \to \mu$, accelerated by N.

Formally, $\bar{Q}_N^+(t) = \frac{1}{N}Q_N^+(t) \xrightarrow{d} \bar{Q}(0)^+$ is a **fluid limit** , since

$$\bar{Q}_N^+(t) = \frac{1}{N}Q_N^+(t)(\textbf{ with } \lambda^+, \mu^+) \stackrel{d}{=} \frac{1}{N}Q_N^+(Nt)(\textbf{ with } \bar{\lambda}, \bar{\mu})$$

The limit is deterministic, hence a degenerate stationary distribution.

To get a **diffusion limit**, **accelerate**:

$$\hat{Q}_N^+(t) = \frac{1}{N}Q_N^+(Nt)(\textbf{ with } \lambda^+, \mu^+) \stackrel{d}{=} \frac{1}{N}Q_N^+(N^2t)(\textbf{ with } \bar{\lambda}, \bar{\mu})$$

Simulations of $\hat{\mathrm{Q}}_{\mathrm{N}}(\mathrm{t}) = \frac{1}{N}[\mathrm{Q}_{\mathrm{N}}(\mathrm{Nt}) - \mathrm{N}]$



Theorem: Let $\lim_{N\to\infty} N(1-\rho_N) = \gamma, \quad 0 < \gamma < \infty.$

Define $\hat{Q}_N(t) = \frac{1}{N}[Q_N(Nt) - N]$. Then

$$\hat{\mathrm{Q}}_{\mathrm{N}}(\mathrm{t}) \overset{\mathrm{d}}{\to} \hat{\mathrm{Q}}(\mathrm{t}) \overset{\mathrm{d}}{=} \mathrm{RBM}(-\gamma\mu, \mathbf{2}\mu).$$

Proof:

Diffusion: $\sigma_N(x) \to \sigma(x) = (2 - x^-)\mu$

Drift: $\mu_N(x) \to \mu(x) = \begin{cases} -\mu\gamma & x \geq 0 \\ +\infty & x \leq 0 \end{cases}$

Via random time-change: $\hat{Q}(t) \overset{d}{=}$ RBM $(-\gamma\mu, 2\mu)$.

(RBM ≥ 0: Since the infinitesimal drift "is $+\infty$" at $x \leq 0$.)

Note: $\hat{Q}(\infty) \overset{d}{=} exp(\gamma)$, establishing convergence over $0 \leq t \leq \infty$.

Note: $V_N(t) \overset{d}{\to} \hat{V}(t) \overset{d}{=} \frac{1}{\mu}\hat{Q}(t), \ 0 \leq t \leq \infty$: no scaling needed.

(Check at $t = \infty$, using Haji & Newell).